

## BENADRYL (DIPHENHYDRAMIN HCL) DOSAGE CHART

GIVE THE WEIGHT APPROPRIATE DOSAGE LISTED BELOW EVERY 4-6 HOURS AS NEEDED

OR AS RECOMMENDED BY YOUR CHILD'S PHYSICIAN



Weight (in pounds)	Elixir (12.5mg/5mL)	FastMelt Chewable (12.5mg tablets)	Tablets/Capsules & Quick Dissolve Strips (25mg tablets or 25 mg strips)
8-9	1.25mL (1/4 tsp)		
10-11	2mL		
12-16	2.5mL (½ tsp)		
17-21	3.75mL (3/4 tsp)		
22-25	4mL		
26-33	5mL (1 tsp)	1 FastMelt	
33-39	6mL	1 FastMelt	
40-45	7mL (1 ½ tsp)	1 ½ FastMelt	
46-52	9mL	1 ½ FastMelt	
53-58	10mL (2 tsp)	2 FastMelt	1 tab/cap/strip
59-65	11mL	2 FastMelt	1 tab/cap/strip
66-72	12mL (2 ½ tsp)	2 ½ FastMelt	1 ½ tabs/caps/strips
73-80	14mL	2 ½ FastMelt	1 ½ tabs/caps/strips
81-88	15mL (3 tsp)	3 FastMelt	1 ½ tabs/caps/strips
89-99	17mL	3 FastMelt	2 tabs/caps/strips
100-109	18mL	4 FastMelt	2 tabs/caps/strips
≥100	20mL (4 tsp)	4 FastMelt	2 tabs/caps/strips

**\*PLEASE NOTE**, this is the maximum recommended safe dosage based on your child's weight. If the listed dosage makes your child too sleepy, you may need to decrease the amount of medicine given in each dose.

## BENADRYL (DIPHENHYDRAMIN HCL) DOSAGE CHART

GIVE THE WEIGHT APPROPRIATE DOSAGE LISTED BELOW EVERY 4-6 HOURS AS NEEDED

OR AS RECOMMENDED BY YOUR CHILD'S PHYSICIAN



Weight (in pounds)	Elixir (12.5mg/5mL)	FastMelt Chewable (12.5mg tablets)	Tablets/Capsules & Quick Dissolve Strips (25mg tablets or 25 mg strips)
8-9	1.25mL (1/4 tsp)		
10-11	2mL		
12-16	2.5mL (½ tsp)		
17-21	3.75mL (3/4 tsp)		
22-25	4mL		
26-33	5mL (1 tsp)	1 FastMelt	
33-39	6mL	1 FastMelt	
40-45	7mL (1 ½ tsp)	1 ½ FastMelt	
46-52	9mL	1 ½ FastMelt	
53-58	10mL (2 tsp)	2 FastMelt	1 tab/cap/strip
59-65	11mL	2 FastMelt	1 tab/cap/strip
66-72	12mL (2 ½ tsp)	2 ½ FastMelt	1 ½ tabs/caps/strips
73-80	14mL	2 ½ FastMelt	1 ½ tabs/caps/strips
81-88	15mL (3 tsp)	3 FastMelt	1 ½ tabs/caps/strips
89-99	17mL	3 FastMelt	2 tabs/caps/strips
100-109	18mL	4 FastMelt	2 tabs/caps/strips
≥100	20mL (4 tsp)	4 FastMelt	2 tabs/caps/strips

**\*PLEASE NOTE**, this is the maximum recommended safe dosage based on your child's weight. If the listed dosage makes your child too sleepy, you may need to decrease the amount of medicine given in each dose.