

Ibuprofen (common brand names: Motrin, Advil)

Give every 6 to 8 hours (as needed for fever or pain) *NOT advised for children under 6 months old*

Weight (in pounds)	Approximate Dose	Infant's Concentrated Drops	Children's Liquid Suspension	Junior Strength Chewable Tablets	Adult Regular Strength Tab
		(50mg/ 1.25 ml)	(100mg/5ml)	(100 mg) each	(200 mg) each
6 to 11	NOT Recommended				
12 to 17	50 mg	1.25 ml	½ tsp = 2.5 ml		
18 to 23	75 mg	1.875 ml	3/4 tsp = 3.75 ml		
24 to 35	100 mg	2.5 ml	1 tsp = 5 ml	1	
36 to 47	150 mg		1.5 tsp = 7.5 ml	1.5	
48 to 59	200 mg		2 tsp = 10 ml	2	1
60 to 71	250 mg		2.5 tsp = 12.5 ml	2.5	1
72 to 95	300 mg		3 tsp = 15 ml	3	1.5
95 and above	400 mg		4 tsp = 20 ml	4	2

Acetaminophen (common brand name: Tylenol) Give every 4 to 6 hours (as needed for fever or pain)



Weight (in pounds)	Approximate Dose	Children's and Infant Liquid Suspension	Children's Chewable Tablets	Junior Strength Chewable Tablets	Adult Regular Strength Tab
		(160mg/5ml)	(80mg) each	(160 mg) each	(325mg) each
6 to 11	40 mg	1/4 tsp = 1.25 ml			
12 to 17	80 mg	½ tsp = 2.5 ml			
18 to 23	120 mg	3/4 tsp = 3.75 ml			
24 to 35	160 mg	1 tsp = 5 ml	2	1	
36 to 47	240 mg	1.5 tsp = 7.5 ml	3	1.5	
48 to 59	320 mg	2 tsp = 10 ml	4	2	1
60 to 71	400 mg	2.5 tsp = 12.5 ml	5	2.5	1
72 to 95	480 mg	3 tsp = 15 ml	6	3	1.5
95 and above	650 mg	4 tsp = 20 ml		4	2